

Melting point 69 – 72°C

Purity (1) Chloride—Heat 2.0 g of Butyl Parahydroxybenzoate with 50 mL of water, allow to stand in ice water for 1 hour with occasional shaking, then add water to make 100 mL, and filter. Perform the test with 25 mL of the filtrate. Prepare the control solution with 0.50 mL of 0.01 mol/L hydrochloric acid VS (not more than 0.035%).

(2) **Sulfate**—Perform the test with 40 mL of the filtrate obtained in Purity (1). Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.024%).

(3) **Heavy metals**—Dissolve 1.0 g of Butyl Parahydroxybenzoate in 25 mL of acetone, add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: to 2.0 mL of Standard Lead Solution add 25 mL of acetone, 2 mL of dilute acetic acid, and water to make 50 mL (not more than 20 ppm).

(4) **Parahydroxybenzoic acid and salicylic acid**—Dissolve 0.50 g of Butyl Parahydroxybenzoate in 30 mL of diethyl ether, shake with 20 mL of a solution of sodium hydrogen carbonate (1 in 100), wash the separated aqueous layer with two 20-mL portions of diethyl ether, shake the aqueous layer with 5 mL of dilute sulfuric acid and 30 mL of diethyl ether, and allow to stand. Shake gently the separated diethyl ether layer with 10 mL of water, remove the aqueous layer after allowing the mixture to stand, filter the diethyl ether, wash the vessel and the filter paper with a small amount of diethyl ether, evaporate the diethyl ether from the combined filtrate and washings on a water bath, and dry the residue in a desiccator (silica gel) to constant mass: the mass of the residue is not more than 5.0 mg. Warm the residue with 5 mL of water, filter, and to the filtrate add 2 to 3 drops of dilute iron (III) chloride TS: no purple color develops.

(5) **Readily carbonizable substances**—Perform the test with 0.50 g of Butyl Parahydroxybenzoate. The solution has no more color than Matching Fluid D.

Loss on drying Not more than 0.5% (2 g, silica gel, 3 hours).

Residue on ignition Not more than 0.10% (1 g).

Assay Weigh accurately about 2 g of Butyl Parahydroxybenzoate, previously dried, add exactly 40 mL of 1 mol/L sodium hydroxide VS, and boil for 30 minutes. Cool, and titrate the excess sodium hydroxide with 0.5 mol/L sulfuric acid VS until the solution shows the same color as that of phosphate buffer solution, pH 6.5, to which the same indicator has been added (indicator: 5 drops of bromothymol blue TS). Perform a blank determination.

$$\begin{aligned} \text{Each mL of 1 mol/L sodium hydroxide VS} \\ = 194.23 \text{ mg of } C_{11}H_{14}O_3 \end{aligned}$$

Containers and storage Containers—Well-closed containers.

Cacao Butter

Oleum Cacao

カカオ脂

Cacao Butter is the fat obtained from the seed of *Theobroma cacao* Linné (*Sterculiaceae*).

Description Cacao Butter occurs as a yellowish white, hard, brittle mass. It has a slight, chocolate-like odor, and has no odor of rancidity.

It is freely soluble in diethyl ether and in petroleum ether, soluble in boiling ethanol (99.5), and very slightly soluble in ethanol (95).

Congealing point of the fatty acids: 45 – 50°C

Melting point 31–35°C (Cram the sample into a capillary tube without melting the sample, then follow Method 2).

Specific gravity d_{20}^{40} : 0.895 – 0.904

Acid value Not more than 3.0.

Saponification value 188 – 195

Iodine value 35 – 43

Containers and storage Containers—Well-closed containers.

Calcium Hydroxide

Slaked Lime

水酸化カルシウム

Ca(OH)₂: 74.09

Calcium Hydroxide contains not less than 90.0% of Ca(OH)₂.

Description Calcium Hydroxide occurs as a white powder. It has a slightly bitter taste.

It is slightly soluble in water, very slightly soluble in boiling water, and practically insoluble in ethanol (95) and in diethyl ether.

It dissolves in dilute acetic acid, in dilute hydrochloric acid and in dilute nitric acid.

It absorbs carbon dioxide from air.

Identification (1) Mix Calcium Hydroxide with 3 to 4 times its mass of water: the mixture is slushy and is alkaline.

(2) Dissolve 1 g of Calcium Hydroxide in 30 mL of dilute acetic acid, and boil. After cooling, neutralize with ammonia TS: the solution responds to the Qualitative tests (2) and (3) for calcium salt.

Purity (1) Acid-insoluble substances—To 5 g of Calcium Hydroxide add 100 mL of water, add hydrochloric acid dropwise with stirring until the solution becomes acidic, and further add 1 mL of hydrochloric acid. Boil this solution for 5 minutes, cool, and filter through a tared glass filter (G4). Wash the residue with boiling water until the last washing ex-

hibits no turbidity upon addition of silver nitrate TS, and dry at 105°C to constant mass: the mass is not more than 25 mg.

(2) Heavy metals—Dissolve 1.0 g of Calcium Hydroxide in 10 mL of dilute hydrochloric acid, evaporate on a water bath to dryness, dissolve the residue in 40 mL of water, and filter. To 20 mL of the filtrate add 2 mL of dilute acetic acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution as follows: evaporate 5 mL of dilute hydrochloric acid on a water bath to dryness, and add 2 mL of dilute acetic acid, 2.0 mL of Standard Lead Solution and water to make 50 mL (not more than 40 ppm).

(3) Magnesium and alkali metals—Dissolve 1.0 g of Calcium Hydroxide in a mixture of 20 mL of water and 10 mL of dilute hydrochloric acid, boil, neutralize with ammonia TS, and precipitate calcium oxalate completely by adding dropwise ammonium oxalate TS. Heat the mixture on a water bath for 1 hour, cool, dilute with water to 100 mL, shake, and filter. To 50 mL of the filtrate add 0.5 mL of sulfuric acid, evaporate to dryness, and ignite at 600°C to constant mass: the mass of the residue does not exceed 24 mg.

(4) Arsenic—Dissolve 0.5 g of Calcium Hydroxide in 5 mL of dilute hydrochloric acid, and perform the test with this solution as the test solution using Apparatus B (not more than 4 ppm).

Assay Weigh accurately about 1 g of Calcium Hydroxide, dissolve by adding 10 mL of dilute hydrochloric acid, and add water to make 100 mL. Measure 10 mL of this solution, add 90 mL of water and 1.5 mL of 8 mol/L potassium hydroxide TS, shake, allow to stand for 3 to 5 minutes, and then add 0.1 g of NN indicator. Titrate immediately with 0.05 mol/L disodium dihydrogen ethylenediamine tetraacetate VS, until the red-purple color of the solution changes to blue.

Each mL of 0.05 mol/L disodium dihydrogen ethylenediamine tetraacetate VS
= 3.7046 mg of Ca(OH)₂

Containers and storage Containers—Tight containers.

Calcium Oxide

Quick Lime

酸化カルシウム

CaO: 56.08

Calcium Oxide, when incinerated, contains not less than 98.0% of CaO.

Description Calcium Oxide occurs as hard, white masses, containing a powder. It is odorless.

It is very slightly soluble in boiling water, and practically insoluble in ethanol (95).

One gram of Calcium Oxide dissolves almost completely in 2500 mL of water.

It slowly absorbs moisture and carbon dioxide from air.

Identification (1) Moisten Calcium Oxide with water: heat is generated and a white powder is obtained. Mix the

powder with about 5 times its mass of water: the mixture is alkaline.

(2) Dissolve 1 g of Calcium Oxide in 20 mL of water by adding a few drops of acetic acid (31): the solution responds to the Qualitative Tests for calcium salt.

Purity (1) Acid-insoluble substances—Disintegrate 5.0 g of Calcium Oxide with a small amount of water, add 100 mL of water, add dropwise hydrochloric acid with stirring until the solution becomes acidic, and further add 1 mL of hydrochloric acid. Boil the solution for 5 minutes, cool, filter through a glass filter (G4), wash the residue with boiling water until no turbidity is produced when silver nitrate TS is added to the last washing, and dry at 105°C to constant mass: the mass of the residue is not more than 10.0 mg.

(2) Carbonate—Disintegrate 1.0 g of Calcium Oxide with a small amount of water, mix thoroughly with 50 mL of water, allow to stand for a while, remove most of the supernatant milky liquid by decantation, and add an excess of dilute hydrochloric acid to the residue: no vigorous effervescence is produced.

(3) Magnesium and alkali metals—Dissolve 1.0 g of Calcium Oxide in 75 mL of water by adding dropwise hydrochloric acid, and further add 1 mL of hydrochloric acid. Boil for 1 to 2 minutes, neutralize with ammonia TS, add dropwise an excess of hot ammonium oxalate TS, heat the mixture on a water bath for 2 hours, cool, add water to make 200 mL, mix thoroughly, and filter. Evaporate 50 mL of the filtrate with 0.5 mL of sulfuric acid to dryness, and heat the residue strongly at 600°C to constant mass: the mass of the residue is not more than 0.015 g.

Loss on ignition Not more than 10.0% (1 g, 900°C, constant mass).

Assay Weigh accurately about 0.7 g of Calcium Oxide, previously incinerated at 900°C to constant mass and cooled in a desiccator (silica gel), and dissolve in 50 mL of water and 8 mL of diluted hydrochloric acid (1 in 3) by heating. Cool, and add water to make exactly 250 mL. Pipet 10 mL of the solution, add 50 mL of water, 2 mL of 8 mol/L potassium hydroxide TS and 0.1 g of NN indicator, and titrate with 0.02 mol/L disodium dihydrogen ethylenediamine tetraacetate VS, until the red-purple color of the solution changes to blue.

Each mL of 0.02 mol/L disodium dihydrogen ethylenediamine tetraacetate VS
= 1.1215 mg of CaO

Containers and storage Containers—Tight containers.

Dibasic Calcium Phosphate

リン酸水素カルシウム

CaHPO₄·2H₂O: 172.09

Dibasic Calcium Phosphate, when dried, contains not less than 98.0% of CaHPO₄: 136.06.

Description Dibasic Calcium Phosphate occurs as a white, crystalline powder. It is colorless and tasteless.