











B. (3-benzoylphenyl)acetic acid,

C. 3-[(1*S*)-1-carboxyethyl]benzoic acid,D. (2*R*)-2-[3-(4-methylbenzoyl)phenyl]propanoic acid,E. (2*R*)-2-(3-benzoylphenyl)propanamide,F. (2*R*)-2-(3-benzoylphenyl)propanenitrile,G. 3-[(1*S*)-1-cyanoethyl]benzoic acid,

H. 3-(cyanomethyl)benzoic acid,




I. (3-benzoylphenyl)ethanenitrile,

J. (2*R*)-2-[3-(2,4-dimethylbenzoyl)phenyl]propanoic acid,K. mixture of (2*R*)-2-[3-(2,3,4-trimethylbenzoyl)phenyl]propanoic acid and (2*R*)-2-[3-(3,4,5-trimethylbenzoyl)phenyl]propanoic acid,L. (2*R*)-2-[3-(2,4,5-trimethylbenzoyl)phenyl]propanoic acid.

01/2008:1755

**KETOROLAC TROMETAMOL**

## Ketorolacum trometamolum

C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>6</sub>  
[74103-07-4]M<sub>r</sub> 376.4

## DEFINITION

2-Amino-2-(hydroxymethyl)propane-1,3-diol (1*S*)-5-benzoyl-2,3-dihydro-1*H*-pyrrolizine-1-carboxylate.*Content*: 98.5 per cent to 101.5 per cent (dried substance).

## CHARACTERS

*Appearance*: white or almost white, crystalline powder.*Solubility*: freely soluble in water and in methanol, slightly soluble in ethanol (96 per cent), practically insoluble in methylene chloride.

## IDENTIFICATION

Infrared absorption spectrophotometry (2.2.24).

*Comparison*: ketorolac trometamol CRS.

## TESTS

**Solution S.** Dissolve 0.75 g in carbon dioxide-free water R and dilute to 25.0 mL with the same solvent.**Appearance of solution.** Solution S is clear (2.2.1).**pH** (2.2.3): 5.7 to 6.7.

Dilute 5 mL of solution S to 15 mL with carbon dioxide-free water R.

**Absorbance** (2.2.25): maximum 0.10, determined at 430 nm for solution S.

**Related substances.** Liquid chromatography (2.2.29). *Protect the solutions from bright light.*

**Solvent mixture:** tetrahydrofuran R, water R (30:70 V/V).

**Test solution.** Dissolve 20 mg of the substance to be examined in the solvent mixture and dilute to 50 mL with the solvent mixture.

**Reference solution (a).** Dilute 1.0 mL of the test solution to 10.0 mL with the solvent mixture. Dilute 1.0 mL of this solution to 100.0 mL with the solvent mixture.

**Reference solution (b).** Dissolve 2 mg of ketorolac trometamol for peak identification CRS (containing impurities A, B, C and D) in 5 mL of the solvent mixture.

**Column:**

- size:  $l = 0.25$  m,  $\varnothing = 4.6$  mm;
- stationary phase: octylsilyl silica gel for chromatography R (5  $\mu$ m);
- temperature: 40 °C.

**Mobile phase:** mix 30 volumes of tetrahydrofuran R with 70 volumes of a solution prepared as follows: dissolve 5.75 g of ammonium dihydrogen phosphate R in 900 mL of water R, adjust to pH 3.0 with phosphoric acid R and dilute to 1000 mL with water R.

**Flow rate:** 1.5 mL/min.

**Detection:** spectrophotometer at 313 nm.

**Injection:** 10  $\mu$ L.

**Run time:** 3 times the retention time of ketorolac.

**Identification of impurities:** use the chromatogram supplied with ketorolac trometamol for peak identification CRS and the chromatogram obtained with reference solution (b) to identify the peaks due to impurities A, B, C and D.

**Relative retention** with reference to ketorolac (retention time = about 10 min): impurity C = about 0.5; impurity A = about 0.6; impurity D = about 0.7; impurity B = about 0.9.

**System suitability:** reference solution (b):

- resolution: minimum 1.5 between the peaks due to impurity B and ketorolac.

**Limits:**

- correction factors: for the calculation of content, multiply the peak areas of the following impurities by the corresponding correction factor: impurity A = 0.67; impurity B = 0.52; impurity C = 2.2;
- impurities A, B, C, D: for each impurity, not more than the area of the principal peak in the chromatogram obtained with reference solution (a) (0.1 per cent);
- unspecified impurities: for each impurity, not more than the area of the principal peak in the chromatogram obtained with reference solution (a) (0.10 per cent);
- total: not more than 10 times the area of the principal peak in the chromatogram obtained with reference solution (a) (1.0 per cent);
- disregard limit: 0.5 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.05 per cent).

**Heavy metals** (2.4.8): maximum 20 ppm.

1.0 g complies with test F. Prepare the reference solution using 2 mL of lead standard solution (10 ppm Pb) R.

**Loss on drying** (2.2.32): maximum 0.5 per cent, determined on 1.000 g by drying *in vacuo* at 60 °C for 3 h.

**Sulfated ash** (2.4.14): maximum 0.1 per cent, determined on 1.0 g.

**ASSAY**

Dissolve 0.300 g in 60 mL of anhydrous acetic acid R. Titrate with 0.1 M perchloric acid, determining the end-point potentiometrically (2.2.20).

1 mL of 0.1 M perchloric acid is equivalent to 37.64 mg of C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>6</sub>.


**STORAGE**

Protected from light.


**IMPURITIES**

**Specified impurities:** A, B, C, D.

**Other detectable impurities** (the following substances would, if present at a sufficient level, be detected by one or other of the tests in the monograph. They are limited by the general acceptance criterion for other/unspecified impurities and/or by the general monograph *Substances for pharmaceutical use* (2034). It is therefore not necessary to identify these impurities for demonstration of compliance. See also 5.10. *Control of impurities in substances for pharmaceutical use*): E, F, G, H, I, J.



- A. R1 = H, R2 = OH: (1RS)-5-benzoyl-2,3-dihydro-1H-pyrrolizin-1-ol,
- B. R1 + R2 = O: 5-benzoyl-2,3-dihydro-1H-pyrrolizin-1-one,
- D. R1 = CO<sub>2</sub>H, R2 = OCH<sub>3</sub>: (1RS)-5-benzoyl-1-methoxy-2,3-dihydro-1H-pyrrolizine-1-carboxylic acid,
- E. R1 = H, R2 = CO-NH-C(CH<sub>2</sub>OH)<sub>2</sub>: (1RS)-5-benzoyl-N-[2-hydroxy-1,1-bis(hydroxymethyl)ethyl]-2,3-dihydro-1H-pyrrolizine-1-carboxamide,
- G. R1 = CO<sub>2</sub>CH<sub>3</sub>, R2 = OH: methyl (1RS)-5-benzoyl-1-hydroxy-2,3-dihydro-1H-pyrrolizine-1-carboxylate,
- H. R1 = H, R2 = CO<sub>2</sub>CH<sub>3</sub>: methyl (1RS)-5-benzoyl-2,3-dihydro-1H-pyrrolizine-1-carboxylate,
- I. R1 = R2 = H: phenyl(2,3-dihydro-1H-pyrrolizin-5-yl)methanone,
- J. R1 = H, R2 = CO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>: ethyl (1RS)-5-benzoyl-2,3-dihydro-1H-pyrrolizine-1-carboxylate,




- C. R6 = CO-C<sub>6</sub>H<sub>5</sub>, R7 = H: (1RS)-6-benzoyl-2,3-dihydro-1H-pyrrolizine-1-carboxylic acid,
- F. R6 = H, R7 = CO-C<sub>6</sub>H<sub>5</sub>: (1RS)-7-benzoyl-2,3-dihydro-1H-pyrrolizine-1-carboxylic acid.

07/2010:1592

## KETOTIFEN HYDROGEN FUMARATE

Ketotifeni hydrogenofumaras



C<sub>23</sub>H<sub>23</sub>NO<sub>5</sub>S  
[34580-14-8]

$M_r$  425.5