

Assay Weigh accurately about 0.5 g of Propranolol Hydrochloride, previously dried, dissolve in 20 mL of acetic acid (100), add 30 mL of acetic anhydride, and titrate with 0.1 mol/L perchloric acid VS (potentiometric titration). Perform a blank determination, and make any necessary correction.


Each mL of 0.1 mol/L perchloric acid VS
= 29.581 mg of $C_{16}H_{21}NO_2 \cdot HCl$

Containers and storage Containers—Well-closed containers.

Storage—Light-resistant.

Propylthiouracil

プロピルチオウラシル



$C_7H_{10}N_2OS$: 170.23
2,3-Dihydro-6-propyl-2-thioxopyrimidin-4(1*H*)-one
[51-52-5]

Propylthiouracil, when dried, contains not less than 98.0% of $C_7H_{10}N_2OS$.

Description Propylthiouracil occurs as a white powder. It is odorless, and has a bitter taste.

It is sparingly soluble in ethanol (95), and very slightly soluble in water and in diethyl ether.

It dissolves in sodium hydroxide TS and in ammonia TS.

Identification (1) Shake well 0.02 g of Propylthiouracil with 7 mL of bromine TS for 1 minute, and heat until the color of bromine TS disappears. Cool, filter, and add 10 mL of barium hydroxide TS to the filtrate: a white precipitate is produced. The color of the precipitate does not turn purple within 1 minute.

(2) To 5 mL of a hot saturated solution of Propylthiouracil add 2 mL of a solution of sodium pentacyanoammine ferroate (II) *n*-hydrate (1 in 100): a green color develops.

Melting point 218 – 221°C

Purity (1) Sulfate—Triturate Propylthiouracil finely in a mortar. To 0.75 g of the powder add 25 mL of water, heat for 10 minutes on a water bath, cool, filter, and wash the residue with water until the volume of the filtrate becomes 30 mL. To 10 mL of the filtrate add 1 mL of dilute hydrochloric acid and water to make 50 mL, and perform the test using this solution as the test solution. Prepare the control solution with 0.40 mL of 0.005 mol/L sulfuric acid VS (not more than 0.077%).

(2) Thiourea—Dissolve 0.30 g of Propylthiouracil in 50 mL of water by heating under a reflux condenser for 5 minutes, cool, and filter. To 10 mL of the filtrate add 3 mL of ammonia TS, shake well, and add 2 mL of silver nitrate TS: the solution has no more color than the following control solution.

Control solution: Weigh exactly 0.060 g of thiourea, and dissolve in water to make exactly 100 mL. Pipet 1 mL of this solution, add water to make exactly 100 mL, and proceed with 10 mL of this solution in the same manner.

Loss on drying Not more than 0.5% (1 g, 105°C, 2 hours).

Residue on ignition Not more than 0.10% (1 g).

Assay Weigh accurately about 0.3 g of Propylthiouracil, previously dried, and add 30 mL of water. Add 30 mL of 0.1 mol/L sodium hydroxide VS from a burette, heat to boil, and dissolve by stirring. Wash down the solid adhering to the wall of the flask with a small amount of water, and add 50 mL of 0.1 mol/L silver nitrate VS with stirring. Boil gently for 5 minutes, add 1 to 2 mL of bromothymol blue TS, and titrate with 0.1 mol/L sodium hydroxide VS until a persistent blue-green color develops. Determine the total volume of 0.1 mol/L sodium hydroxide VS consumed.

Each mL of 0.1 mol/L sodium hydroxide VS
= 8.512 mg of $C_7H_{10}N_2OS$

Containers and storage Containers—Well-closed containers.

Storage—Light-resistant.

Propylthiouracil Tablets

プロピルチオウラシル錠

Propylthiouracil Tablets contain not less than 93% and not more than 107% of the labeled amount of propylthiouracil ($C_7H_{10}N_2OS$: 170.23).

Method of preparation Prepare as directed under Tablets, with Propylthiouracil.

Identification To a quantity of powdered Propylthiouracil Tablets, equivalent to 0.3 g of Propylthiouracil according to the labeled amount, add 5 mL of ammonia TS, allow to stand for 5 minutes with occasional shaking, add 10 mL of water, and centrifuge. To the supernatant liquid add acetic acid (31), collect the precipitate produced, recrystallize from water, and dry at 105°C for 1 hour: it melts between 218°C and 221°C. Proceed with the residue as directed in the Identification under Propylthiouracil.

Dissolution test Perform the test with 1 tablet of Propylthiouracil Tablets at 75 revolutions per minute according to Method 2 under the Dissolution Test, using 900 mL of diluted phosphate buffer solution, pH 6.8, (1 in 2) as the test solution. Take 20 mL or more of the dissolved solution 30 minutes after starting the test, and filter through a membrane filter with pore size of not more than 0.8 μm. Discard the first 10 mL of the filtrate, and use the subsequent as the sample solution. Separately, weigh accurately about 0.05 g of propylthiouracil for assay, previously dried at 105°C for 3 hours, dissolve in diluted phosphate buffer solution, pH 6.8, (1 in 2) to make exactly 1000 mL, and use this solution as the standard solution. Determine the absorbances, A_T and A_S , of the sample solution and the standard solution at 274 nm as directed under the Ultraviolet-visible Spectrophotometry.

The dissolution rate of Propylthiouracil Tablets in 30