

Tolerances: NLT 80% (Q) of the labeled amount of $C_{16}H_{19}N_3O_5S$ is dissolved.

- **UNIFORMITY OF DOSAGE UNITS** (905): Meet the requirements

SPECIFIC TESTS

- **MICROBIAL ENUMERATION TESTS** (61) and **TESTS FOR SPECIFIED MICROORGANISMS** (62): The total aerobic microbial count does not exceed 10^3 cfu/g, and the total combined molds and yeasts count does not exceed 10^2 cfu/g.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, and store at controlled room temperature.
- **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.
- **USP REFERENCE STANDARDS** (11)
USP Amoxicillin RS

Amoxicillin and Clavulanate Potassium for Oral Suspension

DEFINITION

Amoxicillin and Clavulanate Potassium for Oral Suspension contains the equivalent of NLT 90.0% and NMT 120.0% of the labeled amount of amoxicillin ($C_{16}H_{19}N_3O_5S$) and the equivalent of NLT 90.0% and NMT 125.0% of the labeled amount of clavulanic acid ($C_8H_9NO_5$). It contains one or more suitable buffers, colors, flavors, preservatives, stabilizers, sweeteners, and suspending agents.

IDENTIFICATION

- The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the *Assay*.

ASSAY

PROCEDURE

Buffer: 7.8 g of monobasic sodium phosphate in 900 mL of water. Adjust with phosphoric acid or 10 N sodium hydroxide to a pH of 4.4 ± 0.1 , and dilute with water to 1000 mL.
Mobile phase: Methanol and *Buffer* (1:19). Pass through a suitable filter.

Standard solution: 0.5 mg/mL of USP Amoxicillin RS and 0.2 mg/mL of USP Clavulanate Lithium RS in water

Sample solution: Equivalent to 0.5 mg/mL of amoxicillin, from constituted Amoxicillin and Clavulanate Potassium for Oral Suspension in water. Stir by mechanical means for 10 min, and filter. [NOTE—Constitute as directed in the labeling; use within 1 h.]

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 220 nm

Column: 4-mm \times 30-cm; 3- to 10- μ m packing L1

Flow rate: 2 mL/min

Injection size: 20 μ L

System suitability

Sample: *Standard solution*

[NOTE—The relative retention times for clavulanic acid and amoxicillin are about 0.5 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 3.5 between the amoxicillin and clavulanic acid peaks

Tailing factor: NMT 1.5 for each analyte peak

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of $C_{16}H_{19}N_3O_5S$ in the Amoxicillin and Clavulanate Potassium for Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times F \times 100$$

- r_U = peak response of amoxicillin from the *Sample solution*
 - r_S = peak response of amoxicillin from the *Standard solution*
 - C_S = concentration of USP Amoxicillin RS in the *Standard solution* (mg/mL)
 - C_U = nominal concentration of amoxicillin in the *Sample solution* (mg/mL)
 - P = potency of amoxicillin in USP Amoxicillin RS (μ g/mg)
 - F = conversion factor, 0.001 mg/ μ g
- Calculate the percentage of $C_8H_9NO_5$ in the Amoxicillin and Clavulanate Potassium for Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times 100$$

- r_U = peak response of clavulanic acid from the *Sample solution*
- r_S = peak response of clavulanic acid from the *Standard solution*
- C_S = concentration of USP Clavulanate Lithium RS in the *Standard solution* (mg/mL)
- C_U = nominal concentration of clavulanic acid in the *Sample solution* (mg/mL)
- P = potency of clavulanic acid in USP Clavulanate Lithium RS (mg/mg)

Acceptance criteria: 90.0%–120.0% of the labeled amount of $C_{16}H_{19}N_3O_5S$ and 90.0%–125.0% of the labeled amount of $C_8H_9NO_5$

PERFORMANCE TESTS

DELIVERABLE VOLUME (698)

For powder packaged in multiple-unit containers: Meets the requirements

UNIFORMITY OF DOSAGE UNITS (905)

For powder packaged in single-unit containers: Meets the requirements

SPECIFIC TESTS

- **PH** (791): 3.8–6.6, in the suspension constituted as directed in the labeling, the test being performed immediately after constitution
- **MICROBIAL ENUMERATION TESTS** (61) and **TESTS FOR SPECIFIED MICROORGANISMS** (62): The total aerobic microbial count does not exceed 10^2 cfu/g, and the total combined molds and yeasts count does not exceed 5×10^1 cfu/g.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, at controlled room temperature.
- **USP REFERENCE STANDARDS** (11)
USP Amoxicillin RS
USP Clavulanate Lithium RS

Amoxicillin and Clavulanate Potassium Tablets

DEFINITION

Amoxicillin and Clavulanate Potassium Tablets contain the equivalent of NLT 90.0% and NMT 120.0% of the labeled amounts of amoxicillin ($C_{16}H_{19}N_3O_5S$) and clavulanic acid ($C_8H_9NO_5$).

IDENTIFICATION

- The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the *Assay*.

ASSAY

PROCEDURE

Buffer: 7.8 g of monobasic sodium phosphate in 900 mL of water. Adjust with phosphoric acid or 10 N sodium hydroxide to a pH of 4.4 ± 0.1 , and dilute with water to 1000 mL.

Mobile phase: Methanol and Buffer (1:19). Pass through a suitable filter.

Standard solution: 0.5 mg/mL of USP Amoxicillin RS and 0.2 mg/mL of USP Clavulanate Lithium RS in water

Sample stock solution: Dissolve NLT 10 Tablets in water with the aid of mechanical stirring. Transfer to a suitable volumetric flask, and dilute with water to volume.

Sample solution: Dilute a suitable volume of the Sample stock solution filtrate with water to obtain a solution containing 0.5 mg/mL of amoxicillin. [NOTE—Use the Sample solution within 1 h.]

Chromatographic system

(See Chromatography <621>, System Suitability.)

Mode: LC

Detector: UV 220 nm

Column: 4-mm × 30-cm; 3- to 10- μm packing L1

Flow rate: 2 mL/min

Injection size: 20 μL

System suitability

Sample: Standard solution

[NOTE—The relative retention times for clavulanic acid and amoxicillin are 0.5 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 3.5 between the amoxicillin and clavulanic acid peaks

Tailing factor: NMT 1.5 for each analyte peak

Relative standard deviation: NMT 2.0%

Analysis

Samples: Standard solution and Sample solution

Calculate the percentage of C₁₆H₁₉N₃O₅S in each Tablet taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times F \times 100$$

r_U = peak response of amoxicillin from the Sample solution

r_S = peak response of amoxicillin from the Standard solution

C_S = concentration of USP Amoxicillin RS in the Standard solution (mg/mL)

C_U = nominal concentration of amoxicillin in the Sample solution (mg/mL)

P = potency of USP Amoxicillin RS (μg/mg)

F = conversion factor, 0.001 mg/μg

Calculate the percentage of C₈H₉NO₅ in each Tablet taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times 100$$

r_U = peak response of clavulanic acid from the Sample solution

r_S = peak response of clavulanic acid from the Standard solution

C_S = concentration of USP Clavulanate Lithium RS in the Standard solution (mg/mL)

C_U = nominal concentration of clavulanic acid in the Sample solution (mg/mL)

P = potency of clavulanic acid in USP Clavulanate Lithium RS (mg/mg)

Acceptance criteria: 90.0%–120.0%

PERFORMANCE TESTS

- **DISINTEGRATION (701):** Tablets labeled for veterinary use only; 30 min, simulated gastric fluid TS being substituted for water in the test

- **DISSOLUTION (711)**

[NOTE—Tablets labeled for veterinary use only are exempt from this requirement.]

Test 1

Medium: Water; 900 mL

Apparatus 2: 75 rpm

Time: 30 min; or 45 min where the Tablets are labeled as chewable

Analysis: Determine the amount of C₁₆H₁₉N₃O₅S and C₈H₉NO₅ dissolved, using the Analysis set forth in the Assay, making any necessary volumetric adjustments.

Tolerances: NLT 85% (Q) of the labeled amount of C₁₆H₁₉N₃O₅S and NLT 80% (Q) of the labeled amount of C₈H₉NO₅ are dissolved.

For Tablets labeled as chewable: NLT 80% (Q) of the labeled amounts of C₁₆H₁₉N₃O₅S and C₈H₉NO₅ is dissolved in 45 min.

Test 2: If the product complies with this test, the labeling indicates that it meets USP Dissolution Test 2.

Medium, Apparatus 2, and Analysis: Proceed as directed for Test 1.

Times: 45 min for amoxicillin, and 30 min for clavulanic acid

Tolerances: NLT 85% (Q) of the labeled amount of C₁₆H₁₉N₃O₅S and NLT 80% (Q) of the labeled amount of C₈H₉NO₅ are dissolved.

- **UNIFORMITY OF DOSAGE UNITS (905):** Meet the requirements

SPECIFIC TESTS

- **WATER DETERMINATION, Method I (921):**

Tablet Label Claim Amoxicillin (mg/Tablet)	Acceptance Criteria, NMT (%)
≤250	7.5
>250 and ≤500	10.0
>500	11.0

For products labeled as chewable Tablets:

Tablet Label Claim Amoxicillin (mg/Tablet)	Acceptance Criteria, NMT (%)
≤125	6.0
>125	8.0

For Tablets labeled for veterinary use only: NMT 10.0%

- **MICROBIAL ENUMERATION TESTS (61) and TESTS FOR SPECIFIED MICROORGANISMS (62):** The total aerobic microbial count does not exceed 10³ cfu/g, and the total combined molds and yeasts count does not exceed 10² cfu/g.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers.
- **LABELING:** Label chewable Tablets to include the word “chewable” in juxtaposition to the official name. The labeling indicates that chewable Tablets may be chewed before being swallowed or may be swallowed whole. Tablets intended for veterinary use only are so labeled. When more than one Dissolution test is given, the labeling states the Dissolution test used only if Test 1 is not used.
- **USP REFERENCE STANDARDS (11)**
 - USP Amoxicillin RS
 - USP Clavulanate Lithium RS

Amoxicillin Intramammary Infusion

» Amoxicillin Intramammary Infusion is a suspension of Amoxicillin in a suitable vegetable oil vehicle. It contains not less than 90.0 per cent and not more than 120.0 per cent of the labeled amount of amoxicillin (C₁₆H₁₉N₃O₅S). It contains a suitable dispersing agent and preservative.

Packaging and storage—Preserve in well-closed disposable syringes.

Labeling—Label it to indicate that it is intended for veterinary use only.

USP Reference standards (11)—
USP Amoxicillin RS